


KHASKIN, I. G. and BRODSKIY, A. I.

"Isotopic Change of Hydrogen in Contact with Flint," Dokl. AN SSSR, No.6,
21 Oct 50


Some applications of deuterium and of heavy oxygen to the chemistry of silicon. L. O. Kharin. *Doklady Akad. Nauk S.S.S.R.* 83, 139-141 (1952).—As expected from the analogy with the C—H bond, no isotopic exchange was observed between HSiEt_3 , HSiPh_3 , or HSi(OEt)_3 and D_2O , EtOD , or Et_2ND , even on 160 hrs. heating at 118° with solns. of acids or bases in D_2O or EtOD . H being intermediate on the electronegativity scale between C and Si, the polariza-

tions of the bonds are C—H and Si—H, i.e. nucleophilic substitution is favored with Si. Exchange between silanes and proton donors is little probable, as it should be accompanied by a change of the direction of the polarization of the Si—H bond. In silanols, HSiOH , the Si is more electrophilic than in silanes, and nucleophilic exchange in the OH group should be possible. This was confirmed by expts. with EtSiOH and H_2O enriched with O^{18} ; complete exchange took place both without catalyst and with addns. of acids or bases. As an example, EtSiOH was heated with a soln. of NaOH in H_2O with 124 γ excess d., 2.5 hrs. at 100° ; the excess d. of the H_2O became 103 γ , as compared with 88 γ for full exchange. The heavy EtSiOH produced was then heated with light H_2O , 5 hrs. at 100° ; the H_2O showed an excess d. of 21 γ , as compared with 24 γ for complete exchange. With PhSiOH and H_2O^{18} , 40% exchange was found in 1 hr. at 100° . In the exchange of silanols in an alk. medium, the nucleophilic agent is the OH group; in an acid medium, the interaction with the nucleophilic H_2O proceeds by way of the hydrosilanzium ion. In silica gel dried at 400° , both the O of the structural H_2O , and the nonhydroxyl O are exchanged. A sample contg. 3.08% structural H_2O , heated with H_2O^{18} in a sealed tube 30 hrs. at 100° , exchanged 19% of its O. Silica gel entirely free from structural H_2O as a result of prolonged calcination at 1200° , -


exchanged 17% of its O. In the hydrolysis of Si(OEt)_3 , at 78° , with H_2O^{18} (124 γ excess d.), the EtOH was light both in the absence of a catalyst and with addns. of acid or alkali. This decides against the hydrolysis scheme




and in favor of the scheme



On the other hand, in the hydrolysis of Si(OCOMe)_4 with H_2O^{18} (excess d. 124 γ), which takes place violently at the solid-liquid boundary, the H_2O obtained from the AcOH produced had an excess d. of 45-70 γ . This points to a scheme



to the exclusion of the scheme



With respect to the mobility of H, no H—D exchange was observed at 100° in the absence of a catalyst between Si(OCOMe)_4 and AcOD . The exchange does occur in the presence of some AcONa , and, at the same time, there is an exchange of the Ac groups. In 52 hrs. at 100° , with Si(OCOMe)_4 : AcOD : AcONa = 1:2.3:0.1, 92% of all the H of the system was exchanged. In AcOD : AcOD : AcONa = 1:0.85:0.05, in 15 hrs. at 100° , 33% of the H was exchanged. In this case, too, there is also mutual exchange of the Ac

groups. In AcOH contg. 85% γ of D in the OH group, only 6% γ has passed into the Ac group in 7 months at room temp. The mobility of II in $\text{Sn}(\text{OCOMe})_4$ is further confirmed by its condensation with BaH , which takes place in the presence of AcONa only, giving cinnamic acid with a yield of 18% in 13 hrs. at 155° , and 4% in 120 hrs. at 100° . With Na succinate 8-9% phenylsuccinic acid and some cinnamic acid were obtained in 7-10 hrs. at 100° , but no isoprenylsuccinic acid. N. Thon

KLASKIN, I. G.

USSR .

Hydrogen exchange in acetyl silicate and its condensation
 with benzaldehyde I. G. Khasin, V. A. Khasin, N. A. Khasin
 Institute of Chemistry, Academy of Sciences of the USSR, Moscow
 (Received 10.12.77)

It is shown that the rate of hydrogen exchange in acetyl silicate with benzaldehyde increases with increasing temperature and concentration of the aldehyde. The rate of exchange is also increased by the presence of a small amount of water. The exchange is not affected by the presence of a small amount of acetic acid. The exchange is not affected by the presence of a small amount of acetic acid. The exchange is not affected by the presence of a small amount of acetic acid.

12/12/77

KLASHIN I G.

CZECH

Exchange of information

pyridine, base catalyzed

part of ~~the~~ ~~reaction~~

the reaction

the reaction

changes only ΔH° of the

the reaction

the reaction

the reaction

the reaction

the reaction

the reaction

the reaction

Acetyl succinate, $\text{Si}(\text{OAc})_4$, hydrolyzed under similar con-
ditions, indicated formation of AcOH and corresponding HO

deriv. of Si. in this case the

acetyl

Khaskin, I.G.

Chem Abs V48
1-25-54
Organic Chemistry

Mobility of hydrogen in some organosilicon compounds.
I. G. Khaskin. *Zhur. Obshchei Khim.* 23: 327 (1953). No
exchange of H is observed between HSiEt_3 , HSiPh_3 , and
 HSi(OEt)_3 with D from D_2O , EtOD , and DNEt_3 even
on prolonged heating (up to 115°) in presence of acid
(H_2SO_4) or base (NaOH). The electronegativity of
Si being less than that of H the electrophilic type of ex-
change is improbable while nucleophilic reactions are
possible. Thus R_3SiH react nucleophilically with alkalis,
metal amides, alkoxides, etc. The behavior of Si deriva-
tives is readily explainable on this basis. In interconversion of HSi-
 $(\text{OEt})_3$ with EtOH (EtOD) there is an exchange of H for
EtO group. Thus, heating pure HSi(OEt)_3 in sealed tube
with EtOH to 100° for 125 hrs. gave nearly 50% Si(OEt)_4
with liberation of H. HSiCl_3 , b. 32° , was obtained in 44%
yield from dry HCl and Si at 300° . HSiEt_3 (52%, b. 107° ,
d. 0.7301 , from EtMgBr and HSiCl_3) treated with alc.
 KOH yields EtSiOH , b. $154-6^\circ$, d. 0.8597 , while boiling
with aq. 30% alkali gives $\text{Et}_2\text{SiOSiEt}_2$, b. 223° . Similar re-
action with PhMgBr gave 67% Ph_2SiH , b. 180° , m. 36°
(cf. Reynolds, *et al.*, *C.A.* 23, 5470); boiled with alc. KOH
it gave Ph_2SiOH , m. 151° (from ligroine), while hot aq.
30% KOH gave $\text{Ph}_2\text{SiOSiPh}_2$, m. 222° . HSi(OEt)_3 ob-
tained in 43% yield from HSiCl_3 and alc. EtOH , b. 131° ,
d. 0.8752 ; as the reaction mixt. is allowed to stand for
progressively longer periods more Si(OEt)_4 is formed, ac-
counted for by the above exchange reaction. G. M. K.

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000721910010-0
USSR/Organic Chemistry - Synthetic Organic Chemistry, E-2

Abst Journal: Referat Zhur Khimiya, No 19, 1956, 61493

Author: Khaskin, I. G., Yagupol'skiy, L. M., Fialkov, Yu. A., Yakovleva,
V. Ya., Vishnevskaya, G. I.

Institution: ~~Nov~~ M.V. Lomonosov Chem. Phys. Inst. Acad. Sci. USSR

Title: On Preparation of 2-amino-1-p-nitro-phenylethanol

Original

Periodical: Med. prom-st' SSSR, 1955, No 2, 30-32

Abstract: 2-amino-1-p-nitrophenylethanol (I) is obtained by simultaneous
saponification and amination of the acetate of p-nitrophenyl-
chloromethylcarbinol (II) with aqueous-methanol NH_3 . 0.3 mol I
520 ml 26% NH_3 and 500 ml CH_3OH are heated in an autoclave (55° ,
1.5 od m, 1.5 hours with stirring), boiled down in a flask to 1/3
of initial volume, cooled ($40-50^\circ$) acidified with 27 g 80%
 CH_3COOH + 15 ml water. To the solution are added (after removal
of tarry material) 45 ml 40% NaOH ($15-18^\circ$) to an alkaline reac-
tion, I is filtered off, washed with ice water, pressed; yield
82.5% (on the basis of II), MP $133-134^\circ$ (from alcohol).

KHASKIN, L. G.

✓ ~~Methyl dichloroacetate~~
✓ ~~and Yu. A. Piskunov~~
✓ ~~is heated to 40-45° and then is~~
✓ ~~to give CH₃CO₂Me~~
Chem


6

1. Methylenes and ...
2. ...
3. ...
4. ...
5. ...
6. ...
7. ...
8. ...
9. ...
10. ...
11. ...
12. ...
13. ...
14. ...
15. ...
16. ...
17. ...
18. ...
19. ...
20. ...
21. ...
22. ...
23. ...
24. ...
25. ...
26. ...
27. ...
28. ...
29. ...
30. ...
31. ...
32. ...
33. ...
34. ...
35. ...
36. ...
37. ...
38. ...
39. ...
40. ...
41. ...
42. ...
43. ...
44. ...
45. ...
46. ...
47. ...
48. ...
49. ...
50. ...
51. ...
52. ...
53. ...
54. ...
55. ...
56. ...
57. ...
58. ...
59. ...
60. ...
61. ...
62. ...
63. ...
64. ...
65. ...
66. ...
67. ...
68. ...
69. ...
70. ...
71. ...
72. ...
73. ...
74. ...
75. ...
76. ...
77. ...
78. ...
79. ...
80. ...
81. ...
82. ...
83. ...
84. ...
85. ...
86. ...
87. ...
88. ...
89. ...
90. ...
91. ...
92. ...
93. ...
94. ...
95. ...
96. ...
97. ...
98. ...
99. ...
100. ...

Monocarboxylic acid urethides. I. C. Kozlov, L. A. Kozlova, and O. D. Lavrenko. *Chem. Abstr.* 1957, 51, 12140. The urethides are obtained by condensation of monocarboxylic acid with urea in the presence of catalyst.

"APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721910010-0



APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721910010-0"

YAGUPOL'SKIY, L.M.; VISINEVSKAYA, G.O.; YAVORSKIY, D.F.; GRUZ, B.Ye.;
MAKSIMENKO, A.S.; KHASKIN, I.G.; GONSETSKAYA, Ya.V.; KIPRIANOV,
A.I.

Improvement in the method for producing p-nitrophenylchloro-
methylcarbinole. Med.prom. 13 no.3:20-21 Mr '59.

(MIRA 12:5)

1. Institut organicheskoy khimii AN USSR i Kiyevskiy khimiko-
farmatsévticheskiy zavod imeni M.V.Lomonosova.
(METHANOL)

KHASKIN, I.G.; VISHNEVSKAYA, G.I.; LITVINCHUK, O.D.

Preparation of ureides of some monocarboxylic acids. Zhur.prikl.
khim. 33 no.4:986-988 Ap '60. (MIRA 13:8)
(Ureids)

KHASKIN, I.G.

Some applications of chloral in the synthesis of syntomycin.
Ukr. khim. zhur. 26 no.6:740-743 '60. (MIRA 14:1)

1. Khimiko-farmatsevticheskiy zavod im. Lomonosova.
(Syntomycin) (Chloral)

KHASKIN, I.G.; SERGUCHEV, Yu.A.; PROSHKIN, A.A.; VISHNEVSKAYA, G.I.;
YAVORSKIY, D.F.

Production of trichloroacetic acid from tetrachlorethylene. Med.
prom. 15 no.1:39-42 Ja '61. (MIRA 14:1)

1. Institut ispol'zovaniya gaza Akademii nauk USSR.
(ACETIC ACID)

KHASKIN, I.G.

Catalytic activity of silicon and copper in the synthesis of
prussic acid from ammonia and methane. Ukr. khim. zhur. 27 no.2:189-
190 '61. (MIRA 14:3)

1. Institut ispol'zovaniya gaze AN USSR.
(Silicon) (Copper) (Hydrocyanic acid)

KHASKIN, I.G.; LARIONOV, A.V.

Interaction of galenite with natural gas. Ukr. khim. zhur.
28 no.1:118-121 '62. (MIRA 16:8)

1. Institut ispol'zovaniya gaza AN UkrSSR.

VISHNEVSKAYA, G.I.; KHASKIN, I.G.; BUTLEROVSKIY, M.A.; YAGUPOL'SKIY, L.M.;
LITVINCHUK, O.D.; YAKOVLEVA, V.Ya.; CORBUNOVA, A.D.; KIRIYENKO, S.S.

Preparation of syntomycin by dichloroacetylation of
1-p-nitrophenyl-2-aminoethanol. Ukr. khim. zhur. 29 no. 9: 947-950
'63. (MIRA 17:4)

1. Institut organicheskoy khimii AN UkrSSR.

TSYBUL'SKAYA, G.N.; RUDAVSKIY, V.P.; KHASKIN, I.G.

Herbicidal activity of some aromatic derivatives of trichloroacetamide.
Fiziol. rast. 11 no.2:171-174 Mr-Apr '64. (MIRA 17:4)

1. Scientific Research Institute of State Oil and Chemistry
Committee, Kiev.

ACCESSION NR: AP5019677

UR/0064/65/000/008/0577/0578
547.239.23/113.07+547.297.3.07

AUTHORS: Khaskin, I. G.; Vasil'yeva, Z. A.

TITLE: Production of α, α, β -trichloropropionitrile and α, α, β -trichloropropionic acid

SOURCE: Khimicheskaya promyshlennost', no. 8, 1965, 577-578

TOPIC TAGS: chlorination, chlorine organic compound, trichloropropionitrile, trichloropropionic acid

ABSTRACT: The conditions for the synthesis of the herbicides α, α, β -trichloropropionitrile (A), α, α, β -trichloropropionic acid (B), and the sodium salt of B (C) are investigated. The synthesis is based on the chlorination of acrylonitrile with chlorine in the presence of tertiary amines. The yield of A was 40-45%, of B 30-35%, and of C 30-35%. The sodium salt of B is obtained by neutralization of B with sodium hydroxide.

SUBMITTED: 00

ENCL: 00

SUB CODE: 00

NO REF SCV: 001

OTHER: 023

Card

APPROVED FOR RELEASE: 09/17/2001
ACCESSION NR: AP5023548

UR/0220/65/034/004/0715/0719
632.934.1

AUTHOR: Shomova, Ye. A.; Rudavskiy, V. P.; Khaskin, I. G.

TITLE: Fungicidal activity of some aromatic derivatives of trichloroacetamide

SOURCE: Mikrobiologiya, v. 34, no. 4, 1965, 715-719

TOPIC TAGS: fungicide, aromatic compound, fungus, microbiology

ABSTRACT: The action of trichloroacetamide and 19 aromatic derivatives was tested on five phytopathogenic fungi--*Fusarium oxysporum*, *Botrytis cinerea*, *Alternaria solana*, *Aspergillus niger*, and *Phytophthora blanda*. The results show that the activity of the derivatives varies significantly, with some showing high activity and others showing no activity.

ACCESSION NR:--AP5023548

Presumably the fungicidal activity of the unsubstituted trichloroacetamide and its derivatives is due to their ability to form a covalent bond by acylation to the catalytic trichloroacetamide and the amide competes with the radical metabolites, displacing the latter from the amide bonds of cell systems in the fungi. (Fig. 2, art. has available).

ASSOCIATION: none

SUBMITTED: 11Feb64

ENCL: 00

SUB CODE: OC, CC, LS

NO REF SOV: 000

OTHER: 005

Card 2/2

KHABKIN, I.G.; VASIL'YEV, Z.A.

reduction of α,β -trichloropropionitrile and α,α,β -trichloropropionic acid. khim. prom. 41 no.8:577-578 Ag '65.

(MIRA 18:9)

SHOMOVA, Ye.A., RODAVSKIY, V.P.; KHASKIN, I.G.

Fungicidal activity of some aromatic derivatives of trichloroacetamide. Mikrobiologiya 34 no.4:715-719 J1-Ag '65.

(MIRA 18:10)

and to bind the HCl formed, an excess of the initial amine or a tertiary amine over stoichiometric proportions is used. [WA-50; CBE No. 11]

SUB CODE: 07/ SUBM DATE: 05Jun65/

APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000721910010

Card 1/1.

UDC: 547.495.1.07

ACC NR: AP6011992

(A, N)

SOURCE CODE: UR/0326/66/013/005/0906/0910

AUTHOR: Khaskin, I. G.; Stolper, A. L., Tsybul'skaya, G. N.

ORG: Kiev Branch, State All-Union Scientific Research Institute of the Chlorine Industry (Kiyevskiy filial Gosudarstvennogo soyuznogo nauchno-issledovatel'skogo instituta khlornoy promyshlennosti)

TITLE: Herbicidal activity of certain aromatic derivatives of dichloroacetamide

SOURCE: Fiziologiya rasteniy, v. 13, no. 5, 1966, 906-910

TOPIC TAGS: herbicide, aromatic compound, dichloroacetamide, plant physiology, weed killer, dichloride, amide

ABSTRACT: Results of preliminary tests of the physiological activity of a series of aromatic dichloroacetamide derivatives on mono- and di-cotyledonous seeds are reported. Results of treating the seeds with these preparations are shown in the table. Physiological activity depends on chemical structure. Nos. 19-21 were practically inactive and the greatest effects were shown by compounds 1, 9, 10, 15, and 23. Compound no. 1 was most effective against monocots. Compounds no. 2, 6, 7, 15, 17, and 18 were not very selective. The physiological activity of aryldichloroacetamides is due to their antagonism to certain amino acids necessary for the vital activities of the plant.

Card 1/4

UDC: 631.547+632.954

ACC NR: AP6031992

Table 1. Effects of certain N-aryl-dichloroacetamides on germinating seeds of monocotyledonous and dicotyledonous plants

Preparation no.	Name	Chemical formula	Melting point (°C)	Monocots (oats)			Dicots (sugarbeet)		
				Germination % of controls	length, % of controls	% of controls	Germination % of controls	length, % of controls	% of controls
1	2,2-dichloroacetamide		118-119	0	0	0	20.0	61.3	33.0
2	2,2-dichloro-p-acetotoluidide		152-153	87.0	9.7	61.4	74.0	27.8	23.2
3	2,2-dichloro-o-acetotoluidide		131-132	86.0	7.8	27.7	20.0	48.0	83.6
4	2,2-dichloro-m-acetotoluidide		96-99	91.0	35.0	20.6	30.0	21.0	49.0
5	2,2-dichloro-N-benzylacetamide		85.5-86.5	56.0	18.0	31.0	19.0	58.0	77.0
6	2,2-dichloro-p-hydroxyacetanilide		135-137	84.0	26.0	61.7	81.0	36.1	37.1
7	2,2-dichloro-m-hydroxyacetanilide		148-149	83.0	61.0	87.1	88.0	68.8	77.1
8	2,2-dichloro-o-hydroxyacetanilide		132-133	83.0	37.2	61.2	81.0	55.1	56.2
9	2,2-dichloro-p-acetanilide		130-131	0	0	0	3.2	3.0	2.2
10	2,2-dichloro-o-acetanilide		93-94	53.0	7.7	31.0	36.0	36.7	38.4

Card 2/4

ACC NR: AP6031992

11	2,2-dichloro-m-acetanilide		17-78	63.2	5.4	25.2	58.6	33.3	23.1
12	2,2-dichloro-p-acetophenetide		132.5-161.8	44.4	22.4	31.1	46.3	40.4	17.7
13	2,2-dichloro-p-chloroacetanilide		131-137	78.3	8.8	22.9	100.0	22.2	49.8
14	2,2-dichloro-o-chloroacetanilide		103-105	64.0	20.0	50.0	83.3	100.0	81.2
15	2,2-dichloro-m-chloroacetanilide		98-99	45.0	4.5	5.2	61.0	8.1	12.0
16	2,2-dichloro-p-toluoacetanilide		183-184	53.0	18.2	56.0	74.0	70.0	64.4
17	2,2-dichloro-p-dimethylaminoacetanilide		171-172	83.0	17.0	53.0	82.0	5.7	81.0
18	2,2-dichloro-o-nitroacetanilide		78-80	37.0	62.6	49.4	73.0	54.0	61.8
19	2,2-dichloro-p-carboxyacetanilide		212-243	89.0	107.1	102.3	95.0	104.3	54.8
20	2,2-dichloro-o-carboxyacetanilide		178-179	44.0	78.3	82.8	83.0	53.8	48.0
21	2,2-dichloro-m-carboxyacetanilide		218-219	80.0	83.7	77.8	80.0	78.7	98.7
22	2,2-dichloro-δ-acetonaphthalide		163-165.8	98.4	63.4	41.8	104.5	77.7	81.1
23	2,2-dichloroaceto-p-Xylidide		155-156	68.0	8.8	10.7	87.0	43.3	64.8
24	Control	Water	0	96	100	100	95	100	100

Card 3/4

ACC NR: AP6031992

The toxophoric group is a CHCl_2 group in the alpha position in the amide which corresponds to the CH_2NH_2 in amino acids. It is not conclusive, however, that dichloroacetamides behave like enzymes. When iodine is substituted for chlorine in the p-position, substitution capacity is increased but herbicidal activity is decreased. The most effective compound was 2,2-dichloro-p-acetaniside. [WA-50; CBE No. 12]

SUB CODE: 06/ SUBM DATE: 27May65/ ORIG REF: 002/ OTH REF: 002/

Card 1 of 1

ACC NR: AP6029016

SOURCE CODE: UR/0413/66/000/014/0021/0021

INVENTOR: Khaskin, I. G.; Kondratenko, V. I.; Vdovichenko, V. T.

ORG: none

TITLE: Preparation of α -cyanoisopropyl-N-aryl carbamates. Class 12, No. 183733.

SOURCE: Izobret prom obraz tov zn, no. 14, 1966, 21

TOPIC TAGS: dyanoisopropyl aryl carbamate preparation, cyanoisopropyl aryl chloroformate, primary amine, tertiary amine, organic cyanate compound, amine, carbon compound

ABSTRACT: In the proposed method for the preparation of the title compounds, an α -cyanoisopropyl chloroformate is treated with an amine at -10 to 40°C in an inert solvent (toluene or ethyl ether) and the final product is isolated by a known method. To increase the reaction rate and to bind the HCl formed, an excess of the initial amine or a tertiary amine over stoichiometric proportions is used. [WA-50; CBE No. 11]

SUB CODE: 07/ SUBM DATE: 05Jun65/

Card 1/1.

UDC: 547.495.1.07

ACC NR: AM5027778

Monograph

UN/

Kochenov, M. I.; Abramzon, E. I.; Glikin, A. S.; Goloul'nikov, Ye. M.; Kamkhin, YA.,
B.; Khackin, I. N.; Shlayfer, M. L.

Control and measuring automata and devices for automatic lines (Kontrol'no-izmeri-
tel'nyye avtomaty i pribory dlya avtomaticheskikh liniy) Moscow, Izd-vo
"Mashinostroyeniye", 65. 0371 p. illus. 7,600 copies printed.

TOPIC TAGS: automatic control design, automatic control equipment, electric measu-
ring instrument, error measurement

PURPOSE AND COVERAGE: This book deals with constructions and electrical schemes of
automata and devices as planned by the Main Design Office (OKB) of the State Com-
mittee of Machine Building of Gosplan, U.S.S.R. Based on a survey of various control
and measuring apparatus, recommendations are made for selection of a scheme of
measuring and constructing automata and devices, and for an analysis of admissible
boundaries of errors in measuring by automatic control. Principles methods of tes-
ting the precision of control automata are given. This book is recommended for
technical engineers planning and using control and measuring facilities in machine
building. It can also be useful to higher technical school students.

TABLE OF CONTENTS (abridged);

Ch. I. Automata for final control and sorting of parts --5

Card 1/2

UDC: 620.1-52+681.2:621.90.002.5(022)

ACC NR: AM5027778

- Ch. II. Automata and devices for readjusting or blocking of machines --111
- Ch. III. Devices for control monitoring set up in the machines --188
- Ch. IV. Electrical equipment for control and measuring apparatus --275
- Ch. V. Measuring devices -322
- Ch. VI. Permissible errors of measuring with automatic control of dimensions of parts --353
- Ch. VII. Testing precision of work of the control automata --363

SUB CODE: 13 / SUBM DATE: 06May65/

Card 2/2

SHLEYFER, M.L.; ABRAMZON, E.L.; GLIKIN, A.S.; GOLOUL'NIKOV, Ye.M.;
KAMKHIN, Ya.B.; KRUTIK, Ya.B.; KHASKIN, I.N.; KOCHENOV, M.I.,
kand. tekhn. nauk; PODLAZOV, S.S., inzh. red.; SOLOVOV, V.N.,
inzh. red.; VEDMIDSKIY, A.M., kand. tekhn. nauk, dots.

[Control and measurement automatic machines and instruments
for automatic lines]. Kontrol'no-izmeritel'nye avtomaty i
pribory dlia avtomaticheskikh linii. Moskva, Mashinostroenie,
1965. 371 p. (MIRA 18:8)

KHASKIN, I.N.

Final check of cardan bearings in the automatic shop at the First
State Bearing Plant. Stan. 1 instr. 36 no.2:14-20 F '65.
(MIRA 18:3)

KOCHENOV, M.I.; KHASKIN, I.N.

Electric contact measuring instruments with two floating contacts.
Izm.tekh.no.5:18-20 S-O '56. (MLRA 10:2)
(Electric measurements) (Measuring instruments)

S/121/61/000/009/004/006
D040/D113

AUTHORS: Andreyev, V. I., Goloul'nikov, e. M., Ovcharenko, G. I., and
Khaskin, I. N.

TITLE: Raising the level of measurement techniques

PERIODICAL: Stanki i instrument, no. 9, 1961, 33-36

TEXT: The article lists measuring instruments and automatic measuring process control devices being currently produced by the zavod "Kalibr" ("Kalibr" Plant). The following items are mentioned. (1) A profilograph-profile-meter, developed by "Kalibr" in cooperation with Vsesoyuznyy elektrotekhnicheskiiy institut im. V. I. Lenina (All-Union Electrotechnical Institute im. V. I. Lenin). It is the first Soviet instrument for surface roughness measurements in accordance with the international roughness criterion R_a (mean arithmetical deviation of microscopic unevenness from the mean profile line) that will be introduced in the USSR on January 1, 1962. The instrument consists of a post bearing the measuring table and electric drive, an electric measuring unit, and a recorder; all three separate units weigh 80 kg together and are transportable; the system produces 200,000 times

Card 1/3

Raising the level of measurement techniques

S/121/61/000/009/004/006
D040/D113

magnification, and the feeler exerts pressure not above 0.1 g. (2). A feeler type instrument checking roundness of workpieces by measuring induction and producing records by electro-thermic means on a metallized round diagram. It has been designed in cooperation with ENIMS and is also first of its kind in the USSR. (3) Indicator calipers with "cogged-lever" measuring head and dial, eliminating the usual rocking for finding the real diameter of the bore. Calipers for bores up to 18 mm in diameter have a combination of centering and measuring ball points, and calipers for 18-55 mm bores have a rigid centering bridge. Calipers for above 50 mm are pneumatic and universal, i.e. adjustable in a diameters range with the use of a special setting device that is seen in a photograph. Scales of the measuring heads are graduated in 0.001 mm divisions. (4) Levels with 0.01 mm divisions per meter, for measurement of incline on flat and cylindrical surfaces. The levels have a micrometer head for readings and an optic system for zeroing the bubble in the ampoule. (5) Gage blocks of much higher accuracy than previously, produced in accordance with the latest ~~ГОСТ~~ 9038-59 (GOST 9038-59) standard requirements and having a cohesion force of 5.7 kg-f. (6) An automatic machine sorting balls 1-3 mm in diameter with an accuracy to hundredths of one micron. It is based on measurement of electric induc-

Card 2/3

S/121/61/000/009/004/006
D040/D113

Raising the level of measurement techniques


tion and has the pickup and the electronic measuring unit of a "Kalibr-VEI" ("Kalibr VEI") profilograph-profilometer, and an automatic set-up system moving a master ball once in an hour into measuring position for corrections. The machine has been tested at the 4PT3 (4GPZ) plant. A range of such machines will be produced for balls from 3 to 40 mm and from 0.3 to 1 mm in diameter. (7) "Kalibr-MAMI" ("Kalibr-MAMI") measuring and controlling devices for circular grinders with hydraulic drive working with plunge-cut process. They have been produced in cooperation with MAMI, the Moskovskiy avtomekhanicheskii institut (Moscow Automechanical Institute). The "Kalibr-MAMI" have a measurement range of 6-80 mm and make possible grinding of parts with up to 1.2 mm allowance. In test on "3151" and "3161" grinders of the Khar'kov plant they doubled the work rate, and grinding accuracy corresponded 1st class. (8) A series of measuring-controlling devices, designed at the OKB Mosgorsovnarkhoza (OKB of the Moscow City Sovnarkhoz), for automatic transfer lines. Three of such automatics are briefly described and shown in photographs: for internal combustion engine valves, for universal joint bearing rings, and for tractor wheel axles. Photographs are also given of the profilograph-profilometer, the three types of the calipers, the precision level, the ball-sorting automatic, and the "Kalibr-MAMI". There are 11 figures.

Card 3/3

KHASKIN, Khaim Mendelevich; POPOV, G.G., red.; DONNIKOVA, A.A.,
red.izd-va; GRECHISHCHEVA, V.I., tekhn. red.

[Technical and economic justification in the construction
of enterprises of the forest and wood-using industries] Tekh-
niko-ekonomicheskoe obosnovanie stroitel'stva predpriatii
lesnoi fabrichno-zavodskoi promyshlennosti. Moskva, Goslesbum-
izdat, 1962. 98 p. (MIRA 16:4)

(Wood-using industries)
(Industrial plants—Design and construction)



29

Precipitation method for animal (skin) glue. I. S. Khaskin. *Lezhaya Priim.* 2, No. 1/2, 23 (1952). The raw material is cleaned and the glue boiled as usual. The hot glue soln. is filtered then cooled to approx. 25°. At this stage the concn. of glue in the soln. is 5-7%. To this soln., placed in a wooden tank, is added 1 ltr. vol. of a 28% H_2SO_4 soln. The pptd. gelatin is placed on wooden lattice work covered with sucking and left to drain for 1-2 hrs. It is then pressed to remove more moisture. The gelatin is then melted on a water bath and cast into pieces 15-20 mm. thick. At this stage the gelatin contains not over 30% H_2O . By more exposure to air and without any special drying it will lose another 12-15% of H_2O . The liquor after the gelatin is removed is 15-17% H_2O . It is valuable as such for the leather industry. If desired it can be concd. to 28% H_2SO_4 for reuse. The viscosity of the glue prepd. by the above method was 1.5°, and at times it reached 8-9°, Engler. Its ash content was not over 1% and in a few cases only 1.3%. The sulfate content is 25-30%. It is readily reduced to 6-10% if the dry gelatin is soaked up to 24 hrs. instead of the usual 12 hrs. This process requires no special installations used in glue processing plants. It permits processing of the raw material at or near its source. M. Hosh

KHASKIN, L.S.

Brush production in Germany. Leg.prom. 7 no.8:32-3 of cover. Ag '47.
(MIRA 6:11)

(Germany--Brooms and brushes) (Brooms and brushes--Germany)

FELESHIN, L. S.

TECHNOLOGY

(Obtaining fats from raw material and waste products from the tanning and fur industry). Moskva, Girslegprom. 1951.

Monthly List of Russian Accessions. Library of Congress, November 1952. UNCLASSIFIED.

MOSKALEV, V.M.; KHASKIN, L.S., redaktor; KORNEYEVA, V.I., tekhnicheskii redaktor.

[Textile materials used in the chemical industry] Tekstil'nye materialy, primenyaemye v khimicheskoi promyshlennosti. Moskva, Gos. nauchno-tekhn. izd-vo khim. lit-ry, 1954. 116 p. (MIRA 8:1)
(Chemical industries) (Textile fibers)

KIVMAN, G.Ya., kandidat meditsinskikh nauk; KHASKIN, L.S.

Utilization and sterilization of side products obtained during
production of antibiotics; review of foreign periodical literature.
Antibiotiki 8 no.2:25-36 '55. (MLRA 8:5)

(ANTIBIOTICS, preparation of,
use of side products, review)

(DRUG INDUSTRY,
use of side products in antibiotic indust., review)

KHASKIN, L.

**Efficient utilization of waste leather. (From: Leather Trades Review,
v.117 no.3615, '55). Leg.prem.15[i.e.16] no.3:56 Mr '56.(MLRA 9:7)
(Leather industry)**

NP AF000773

None

00

ENCL 00

EE

000

OTHER 000

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
PROCESSES AND PROPERTIES INDEX																			
<p>KLHASKIN, S.A.</p> <p>CA</p> <p>The utilization of ferrous sulfate from pickling waste waters. S. A. Klhaskin. <i>Vodoumasheniye Soud. 1944, 1939, No. 11-12, 69-72; Khim. Referat. Zhur. 1940, No. 5, 90-7.</i>—The acid waste waters contain H_2SO_4 10-50 and $FeSO_4$ 100-250 g./l. Fe scrap is added to react with the H_2SO_4, the soln. is passed to a Robert vacuum app. (temp. about 60°) with a countercurrent barometric cascade-type condenser. The cooled soln. is passed to crystallizers cooled with circulating salt brine. W. R. Henn</p>										<p>9</p>									
<p>ASAC, S.A. METALLURGICAL LITERATURE CLASSIFICATION</p>										<p>SECOND ORDER</p>									

Subject : USSR/Mining

AID P - 2093

Card 1/1 Pub. 78 - 6/24

Authors : Mongayt, I. L., Konobeyev, S. I. and Khaskin, S. A.

Title : New method in planning oil traps

Periodical: Neft. khoz., v.33, no.4, 28-34, Ap 1955

Abstract : Formulae are given for sediment oil trap tanks or sumps to determine their proper dimensions as dependent upon the specific weights of oil and water, oil concentration, concentration of suspended solids, etc. Charts.

Institution: VNIIVODGEO (All-Union Scientific Research Institute for Water Supply, Sewer Systems, Hydraulic Structures, and Hydrogeological Engineering); AzNII (Azerbaydzhan Scientific Research Institute); UPNII (Ufa Scientific Research Institute)

Submitted : No date

KHASKIN, S.A., inzh. (Moskva)

~~Principles~~ in the construction of water purifying systems in
modern petroleum refineries. Stroi. pred. neft. prom. 3 no.1:10-13
Ja '58. (MIRA 11:3)
(Water--Purification)

KHASKIN, S.A.; VOLKOVA, V.A.

Clarifying reservoirs for waste waters containing petroleum. Vod.
i san. tekhn. no.5:29-31 Ky '58. (MIRA 11:6)
(Sewage--Purification)

KHASKIN, S.A.

Scraping installation for horizontal settling tanks. Vod. i san.
tekh. no.6:13-14 Je '59. (MIRA 12:8)
(Water--Purification)

ZAK, Genrikh Lazarevich, kand.tekhn.nauk; KHASKIN, S.A., red.; OTOCHEVA,
M.A., red.izd-va; SHLIKHT, A.A., tekhn.red.

[Self-purification of water reservoirs; principles underlying
the regulation of hydrological and sanitary-engineering calcula-
tions] Samoochishchenie vodoemov; osnovy ratsionalizatsii gidro-
logicheskikh i sanitarno-tekhnicheskikh raschetov. Moskva, Izd-vo
M-va kommun.khoz.RSFSR, 1960. 159. (MIRA 13:5)
(Water--Purification)

KHASKIN, S.A.

Industrial water supply and sewerage in modern petroleum
refineries. Vod.i san.tekh. no.8:22-24 Ag '60.
(MIRA 13:7)

(Water supply, Industrial) (Sewerage)
(Petroleum refineries--Equipment and supplies)

KHASKIN, S.A.

Purification waste waters from the production of synthetic fatty acids. Zhur. VKHO 6 no.2:188-193 '61.

(Sewage---Purification) (Acids, Fatty)

BALASHOV, A.I.; ARONOV, S.N.; YERESNOV, N.V.; MOSKVITIN, A.S.;
NEMIROVSKIY, D.B. [deceased]; RUBINSHTEYN, S.L.;
POPOVA, V.V.; KHASKIN, S.A.

"Handbook on water supply and sewerage." Reviewed by
A.I. Balashov and others. Vod. i san. tekhn. no.12:32-34
D '62. (MIRA 15:12)

(Water supply)
(Sewerage)

BEKKER, Semen Mikhaylovich, prof.; KHASKIN, Semen Grigor'evich, prof.;
AL'TPOV, V.I., red.; KHARASH, G.A., tekhn. red.

[Women's clinic] Zhenskaia konsul'tatsiia. Leningrad, Medgiz,
1961. 149 p. (MIRA 15:1)
(GENERATIVE ORGANS, FEMALE—DISEASES)
(PREGNANCY, COMPLICATIONS OF)

BELYAYEV, Ye.I., prof. [deceased]; BADYUK, Ye.Ye.; BOGOROV, I.I.,
prof.; BUBLICHENKO, L.I., prof.[deceased]; IL'IN, I.V.,
dots.; KEYLIN, S.L., prof.; MAZHBITS, A.M., prof.;
MALININ, A.I., zasl. deyatel' Kaz.SSR, prof.; MOSHKOV, B.N.,
prof.; NIKOLAYEV, A.P., prof.; PERSIANINOV, L.S., prof.;
POKROVSKIY, V.A., prof.; POLYAKOVA, G.P., kand. med. nauk;
RAFAL'KES, S.B., dots.; KHASKIN, S.G., prof.; SHTERN, I.A.,
prof.

[Multivolume manual on obstetrics and gynecology] Mnogo-
tomnoe rukovodstvo po akusherstvu i ginekologii. Moskva,
Meditsina. Vol.3. Book 2. [Pathology of the labor and post-
natal period. Physiology and pathology of the newborn infant]
Patologiya rodov i poslerodovogo perioda. Fiziologiya i pa-
tologiya novorozhdenno. Pt.1.[Pathology of labor] Patolo-
giya rodov. 1964. 895 p. (MIRA 17:7)

1. Chlen-korrespondent AMN SSSR (for Persianinov). 2. Deystvi-
tel'nyy chlen AMN SSSR (for Nikolayev).

KHASKIN, S.G., prof.

Prevention of suppurative infection in puerperants and newborn infants with staphylococcal anatoxin. Akush. i gin. 20 no.1: 13-17 Ja-F '64. (MIRA 17:8)

1. 2-ya akusherskiye otdeleniye (zav. - prof. S.G. Khaskin)
Instituta akusherstva i ginekologii (dir. - prof. M.A. Petrov-Maslakov) AMN SSSR, Leningrad.

BARTEL'S, A. V., dotsent; RAFAL'KES, S. B., dotsent; KHASKIN, S. G., prof.

Prevention and treatment of lactation mastitis. Akush. i gin.
no.2:3-25 '62. (MIRA 15:6)

(BREAST—DISEASES) (LACTATION)

KHASKIN, V.

Method of calculating the results of the financial activity of
automotive transport enterprises. Avt.transp. 34 no.9:7-8 S '56.
(MLRA 9:11)

1. Zamestitel' direktora avtobazy "Odestorgtrans".
(Transportation, Automotive--Accounting)

MAKHIN'KO, V.I.; KHASKIN, V.V.; SHUL'MAN, G.Ye.

Some features of nitrogen metabolism at a great age. Uch.zap.KHGU
68:193-213 '56 (MIRA 11:11)

1. Kafedra fiziologii cheloveka i zivotnykh Nauchno-issledovatel'-
skogo instituta biologii i biologicheskogo fakul'teta Khar'kovskogo
ordena trudovogo krasnogo znameni gosudarstvennogo universiteta imeni
A.M. Gor'kogo.

(NITROGEN METABOLISM) (OLD AGE)

KHASKIN, V.V.

Physiological effects of temperature on young ducks. Ptitsevodstvo
8 no.12:18-21 D '58. (MIRA 11:12)

1. Ukrainskaya opytnaya stantsiya ptitsevodstva.
(Ducks) (Temperature--Physiological effect)

KHASKIN, V.V.; TITSKIY, I.Ya.

Mixed silage for poultry. Ptitsevodstvo 9 no.8:7-11
Ag '59. (MIRA 12:12)

1. Ukrainskaya opytnaya stantsiya ptitsevodstva.
(Poultry--Feeding and feeds) (Ensilage)

KHASKIN, V.V.

Development of thermoregulation in the domestic duck. Fiziol. zhur.
46 no.12:1489-1496 D '60. (MIRA 14:1)

1. Ukrainskiy nauchno-issledovatel'skiy institut ptitsevodstva,
Khar'kov.

(BODY TEMPERATURE—REGULATION) (DUCKS)

KHASKIN, V.V.

Heat exchange of bird eggs during incubation. Biofizika 6
no. 1:91-99 '61. (MIRA 14:2)

1. Ukrainskiy nauchno-issledovatel'skiy institut ptitsevodstva,
Khar'kov.

(EMBRYOLOGY—BIRDS) (ANIMAL HEAT)

L 19438-63

ACCESSION NR: AP3007181

S/0239/63/049/009/1120/1121

AUTHOR: Khaakin, V. V.

TITLE: A device for the study of gaseous exchange in small animals

SOURCE: Fiziologicheskiy zhurnal SSSR, v. 49, no. 9, 1963, 1120-1121

TOPIC TAGS: oxygen consumption measurement, respirometer, closed circulation respirometer, respiration measurement, animal oxygen consumption rate

ABSTRACT: A device intended for measurement of oxygen-consumption rates of small animals (such as chickens and mice) at different temperatures is described. The machine belongs to the class of respirometers of the closed-circulation type and has the following components (numbers refer to Fig. 1 of the Enclosure): glass animal container 1 (volume, 1 liter) tightly closed with a rubber stopper, two glass tubes and attached rubber hoses 3 and 4 which are connected with CO₂ tank 16; thermometer 5, cross pipe with a

Card 1/3

L 19438-63

ACCESSION NR: AP3007181

three-way gage 6, gasometric buret 7, and manometer 8. During experiments the animal container is immersed in water which fills glass jar 9. To prevent floating, the rubber stopper is firmly attached to a stand, while tubes 11 and 12 are connected with the U-8 ultrathermostat. The oxygen supply is controlled automatically. Container 1 is connected with buret 7 by means of hoses 12, 13, and 14. CO₂ tank 16 is attached to T-frame 17, which in turn is attached to horizontal beam 19. The device is rocked by an electric source to ensure 500 ml of concentrated KOH in the CO₂ tank with the incoming gas. Orig. art. has: 1 figure.

ASSOCIATION: Ukrainskiy nauchno-issledovatel'skiy institut ptitsevodstva, Khar'kov (Ukrainian Scientific Research Institute of Poultry Breeding)

SUBMITTED: 20Aug62

DATE ACQ: 30Sep63

ENCL: 01

SUB CODE: AM

NO REF SOV: 002

OTHER: 000

Card 2/3

L 19438-53

ACCESSION NR: AP3007181

ENCLOSURE: 01




Fig. 1. Device for measuring oxygen-consumption rates of small animals

Card 3/3

TUKALO, Ye.A. [Tukalo, I.E.A.]; KHORON'KO, A.T.; MURATOVA, I.O.; KHASKIN,
Ye.A. [Khaskin, I.E.A.]

Production training for students. Farmutsev. zhur. 17 no.5:82-84
'62. (MIRA 17:9)

1. Kafedra tekhnologii lekarstv Dnepropetrovskogo meditsinskogo
instituta.

FEDOROVSKAYA, N.P.; KHASKINA, I.M.

Micromethod for the determination of chlorine and bromine.
Trudy IGI 21:190-196 '63. (MIRA 16:11)

FEDOROVSKAYA, N.P.; KHASKINA, I.M.; CHUMACHENKO, M.N.

Micromethod for the determination of iodine content.

Trudy IGI 21:197-201 '63.

(MIRA 16:11)

PRILEZHAYEVA, B.N.; FEDOROVSKAYA, N.P.; MIYESSEROVA, L.V.;
DOMANINA, O.N.; KHASKINA, I.M.

Methods of determining varieties of organic sulfur in solid
fuels. Trudy IGI 21:159-168 '63.

Determining sulfur ether in solid fuel by the methyl iodide
method. 202-210 (MIRA 16:11)

FEDOROVSKAYA, N.P.; KHASKINA, I.M.; CHUMACHENKO, M.N.

Simultaneous determination of halides and mercury in halogenated
and mercurated solid fuels. Trudy IGI 8:213-220 '59.

(MIRA 13:1)

(Coal--Analysis)

L 16060-66 FWT(1) GW

ACC NR: AP6004201

SOURCE CODE: UR/0050/66/000/002/0039/0041

AUTHOR: Khaskina, M. I. 268

ORG: Hydrometeorological Scientific-Research Center, SSSR (Gidrometeorologicheskii nauchno-issledovatel'skiy tsentr, SSSR)

TITLE: Prediction of maximal outflows of water¹² at flood stage of a large river according to discharge of small rivers (on the example of the Dnieper near Kiev)

SOURCE: Meteorologiya i gidrologiya, no. 2, 1966, 39-41

TOPIC TAGS: water, hydrology, river, flow measurement, flood

ABSTRACT: A means for predicting maximal outflows of the Dnieper River near Kiev is presented. The method is based upon computation of flood stage hydrographs according to the discharge of smaller rivers. Flows in this river network are given by the formula

$$q = \frac{F}{8} \sum_{i=1}^8 \frac{Q_i}{f_i}$$

where F is the watershed area above Kiev, equal to 320 000 km²; Q_i and f_i are the discharges and areas of each of the eight basins of the smaller rivers.

Card 1/2

UDC: 551.582.215.1 2

L 16060-66
ACC NR: AP6004201

Instantaneous flow in a closed (control) area at a time t is given by the formula

$$Q_t = \sum_{\tau=1}^{\tau_{max}} q_{t-\tau} R(\tau),$$

where $R(\tau)$ is the riverbed runoff curve (effect function), and τ is the runoff time. For the stated problem conditions this equation takes the form

$$\begin{aligned} Q_t = & 0.04 q_{t-4} + 0.08 q_{t-8} + 0.13 q_{t-12} + 0.15 q_{t-16} + \\ & + 0.12 q_{t-20} + 0.09 q_{t-24} + 0.07 q_{t-28} + 0.06 q_{t-32} + \\ & + 0.06 q_{t-36} + 0.05 q_{t-40} + 0.05 q_{t-44} + 0.04 q_{t-48} + \\ & + 0.03 q_{t-52} + 0.02 q_{t-56} + 0.01 q_{t-60}. \end{aligned}$$


Runoff records for the years 1931, 1936-39, and 1945-64 are available for use as inputs to the equation for instantaneous flow. These data are plotted and used in deriving an empirical formula for the time interval for maximum river surge. The proper interpretation of the prediction method is discussed, and the accuracy of the system is evaluated. Use of the method on past occasions resulted in accurate predictions. Orig. art. has: 4 equations and 2 figures.

SUB CODE: 08/ SUBM DATE: 16Jul65/ ORIG REF: 002
Card 2/2

KHASKINA, M.I.

Forecasting the runoff of high water in the Dniaper River near Kiev
based on the runoff of small rivers. Trudy TSIP no.117:87-97 '63.
(MIRA 16:7)

(Dniaper River--Runoff)



KHAZAKI, Ye. Ye.

1/2

Chemical Abst.
Vol. 48 No. 8
Apr. 25, 1954
Organic Chemistry

Sulfonation and sulfonic acids of acidophobic substances.

XXII. Sulfonation of vinyl ethers. A. P. Terent'ev, A. N. Kozlov, B. M. Yurkevich, and E. E. Khachikyan (Moscow State Univ.). *Zhur. Obshch. Khim.* 29, 745-54 (1955); cf. C.A. 48, 3130d. Heating 5 g. 25% soln. of $\text{CH}_2=\text{CHCl}$ in $(\text{CH}_2\text{Cl})_2$ and 0 g. pyridine- SO_3 in an ampul 1.5-2 hrs. at 170° , soln. in H_2O , neutralization with BaCO_3 , steam distn. of the pyridine, filtration of the residue, treatment of the filtrate with C, concn., addn. of MeOH , and extrn. of the product with hot EtOH for 3 days gave 5% $(\text{HCOCH}_2\text{SO}_3)_2\text{Ba}$ (I), formed apparently from the primary product

$\text{CH}_2=\text{CHCl.O.SO}_3\text{O.Ba.O.SO}_3$. Similar reaction of $\text{Me}_2\text{C}=\text{CHBr}$ gave after 6 hrs. at 110° 85% Ba salt of 2-sulfonobutyraldehyde (II), which reduces ammoniacal AgNO_3 ; Pb salt, sirup; Ag salt, insol. in H_2O . $\text{CH}_2=\text{CHOBu}$ (2 g.), 3.2 g. pyridine- SO_3 and 6 ml. $(\text{CH}_2\text{Cl})_2$ heated 9 hrs. at $70-90^\circ$ and treated as above, gave 30% I, forming a monohydrate on crystn. from H_2O ; refluxing the reactants 14 hrs. gives a 32.5% yield; use of dioxane- SO_3 gives 43%. I with S-2-naphthylthiuronium chloride gave the S-2-naphthylthiuronium salt, m. $202-4^\circ$ (from C_6H_6). $\text{CH}_2=\text{CHOAc}$ and pyridine- SO_3 in $(\text{CH}_2\text{Cl})_2$ gave, after 8 hrs. at 120° and the usual treatment, 85% I; dioxane- SO_3 gave 62%. For better isolation of the product and removal of AcOH the product is best refluxed 4 hrs. with 0.2N H_2SO_4 before treatment with BaCO_3 . $\text{H}_3\text{C}=\text{CMeOAc}$ (1 g.) added to 1 g. SO_3 , 6 ml. $(\text{CH}_2\text{Cl})_2$, and 0.9 g. dioxane with ice cooling gave 67% Ba acetonesulfonate monohydrate. Refluxing 28.5 g. iso- PrCHO , 61 g. Ac_2O and 6 g. KOAc 10 hrs. gave 30.5 g. $\text{Me}_2\text{C}=\text{CHOAc}$, b.p. $121-4^\circ$, n_D^{20} 1.4100, which, heated with pyridine- SO_3 10 hrs. at 150° in an ampul, gave 35% II. $\text{CH}_2=\text{CH}_2\text{CHO}$ heated with SO_3 in $(\text{CH}_2\text{Cl})_2$ 12 hrs. on a steam bath, treated with H_2O , freed of Hg salts with H_2S , and neutralized with BaCO_3 , gave 41% Ba sulfacetate monohydrate (from H_2O). Addn. of dioxane dibromide (62 g.) to 21.2 g. $\text{CH}_2=\text{CHOAc}$ with cooling gave 50% $\text{Br-CH}_2\text{CH}_2\text{CH}_2\text{OAc}$, b.p. $101-3^\circ$, n_D^{20} 1.5457, d.n. 1.0170, which (2.5 g.) reduced 1 hr. with 3 g. Na_2SO_3 in 25 ml. H_2O , concn., treated with BaCO_3 , filtered, evaporated, and treated with S-2-

(over)

Sub. Org. Chem.
in. Zeluskiy

KHASKINA-MUNDER, G.N.

EXCERPTA MEDICA Sec.4 Vol.11/4 Med.Microb. etc. April 58

838. LABORATORY DIAGNOSIS OF DOUBTFUL CASES OF SCARLET FEVER
I. (Russian text) - Khaskina-Munder G.N., Biumberg F.M.,
Smirnova E.I. and Elina M.Y. - NAUCH. TRUD. MOSK. INST.
VAKT. SYVOR. 1956, 6 (143-150)

The authors evaluated various laboratory methods of investigation in the diagnosis of doubtful cases of scarlet fever. The methods used gave the following results: (1) Barannikov-Doehle bodies were found in the blood in 16% of the patients under investigation, but were also encountered in other infectious diseases; (2) examination of the throat for haemolytic streptococci was positive in 81% of the patients with clinically obvious infection, and in 14 (36.8%) of 38 patients in whom the diagnosis of scarlet fever had been rejected; (3) cutaneous tests with different doses of streptococcal toxin produced a reaction on the 3rd-5th day in 43.9% of the patients with a doubtful diagnosis; on repeated examination it was found that the majority of cases (28 out of 32) showed a reaction on the 4th-7th day; (4) an increasing agglutinin titre (streptococcus) was observed by the 6-10th day of illness in 89.6% of the patients with uncertain diagnosis; (5) the opsonin-phagocytic reaction proved useless for diagnosis. The authors consider the following tests to be of diagnostic value: bacteriological examination of nose and throat swabs, skin test with streptococcal toxin and determination of the blood agglutinin titre. (S)

KNACKED, H.D.

"Ploskaya Zadacha o Kolebaniyakh Plastiuki ina Poverkhnosti Tyazheloi
Zhidkosti," Izvestiya Akademii Nauk SSSR, Otdel. Tekhnicheskikh Nauk
1982 vyp. 7/8 str. 75 - 94.

KHASKIND, M.D.

Ploskaia zadacha ob ustanovivshikhsia kolebaniakh kryla pod poverkhnost'iu tiazheloi zhidkosti konechnoi glubiny. (Akademiia Nauk SSSR. Izvestiia. Otdelenie tekhnicheskikh nauk, 1942, no. 11-12, p.66-86)

Title tr.: Plane problem of steady oscillations of a wing immersed in heavy fluid of finite depth.

AS262.A6244 1942

SO: Aeronautical Sciences and Aviation in the Soviet Union, Library of Congress, 1955

KR. S. LEB, V. D.

Plaskaya zadacha o glissirovani po poverkhnosti tiazhelei zhidkosti konec noi glubiny. (Akademiia nauk SSSR. Izvestiia. Otdelanie tekhnicheskikh nauk. 1943, no. 1-2, p. 87-90)

Title tr.: Plane problem of planing on the surface of a heavy fluid of finite depth.

AS267.A6344 1943

SO: Aeronautical Sciences and Aviation in the Soviet Union, Library of Congress, 1951

KHASKIND, M.D.

"V Kolebaniya Sistemy Plastinok Na Poverkhnosti Tverdogo Tela,"
Iz Vvedeniya Matematika i Mekhanika Tom 7 vyp. 1 str. 441 - 444, 1961.

MAE. IIL, A. D.

Plaskina zadacha o kolebaniakh tela pod povorkhnost'iu tiazhelei zhidkosti konechnoi glubiny. (Prikladnaya aerostatika i rekonstruktsiya, 1944, v. 6, no. 1, p. 27-300) Summary in English.

Title tr.: Plane problem of oscillations of a body below the surface of a heavy fluid of finite depth.

GA601.F7 1944

SA: Aeronautical Sciences and Aviation in the Soviet Union, Library of Congress, 1955

KHASKIND, M. D.

KHASKIND, M. D.

Ploskaia zadacha o kolebaniakh tela pod poverkhanost'iu tiazheloi zhidkosti konechnoi glubiny. (Prikladnaia matematika i mekhanika, 1944, v. 8, no. 4, p. 287-300)

Summary in English.

Title tr.: Plane problem of oscillations of a body below the surface of a heavy fluid of finite depth.

QA801.P7 1944

SO: Aeronautical Sciences and Aviation in the Soviet Union, Library of Congress, 1955

KHASKIND, M. D.

KHASKIND, M. D.

O postupatel'nom dvizhenii tel pod svobodnoi poverkhnost'iu tiazheloi zhidkosti konechnoi glubiny. (Prikladnaia matematika i mekhanika, 1945, v. 9, no. 1, p. 67-78)

Summary in English.

Bibliography: p. 78.

Title tr.: Translation of bodies below the free surface of a heavy fluid of finite depth.

QA801.P7 1945

SO: Aeronautical Sciences and Aviation in the Soviet Union, Library of Congress, 1955

W. WILD, MT.

"Kolebaniya Sistem Plastinok Na Poverkhnosti Tyazhelei Zhidkosti,"
Prikladnaya Matematika i Mekhanika V. 9, No. 6, str. ... 1945.

Hasland, M. D. Wave resistance of a solid moving
through a fluid of finite depth. *Phil. Mag.* 1954, 9, 1-14.
Also, *Proc. Roy. Soc. London* 1954, 227, 1-14.

This paper deals with the wave resistance of a solid moving
through a fluid of finite depth. The problem is solved for
a solid of arbitrary shape moving with constant velocity
in a fluid of finite depth. The results are expressed in
terms of the wave number k and the velocity V . The
wave resistance is shown to be a function of the wave
number k and the velocity V . The results are
expressed in terms of the wave number k and the velocity
 V . The wave resistance is shown to be a function of
the wave number k and the velocity V .

1 FH

KILGORE, M.D.

Kachka Korablya Na Spokoynoy Vode," Izvestiya Akademii Nauk SSSR, Otdel.
Tekhnicheskikh Nauk 1946, vyp. 1 str. 23 - 24.

KHASKIND, M. D.

"Hydrodynamic Theory of Oscillations of a Ship in Waves," Prik. mat. i mekh.,
10, No.1, 1946 p. 33-66

Hashing, M. D. Waves are re-
in shall w water. The
SSNR (Peak Mean Value) is

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

1.0 (dB)

PA40793

USSR/Physics
Fluids, Compressible
Acoustic Radiators

Aug 1946

"Acoustical Radiation of Oscillating Bodies in a
Compressed Liquid," M. D. Khaskind, 13 pp

"Zhur Eksp 1 Teoret Fiz" Vol XVI, No 7

Theoretical investigation of the problem of acoustical radiation of an oscillating body in a compressed liquid, more specifically, the harmonic oscillation of a solid and deformable body, has been carried out. During oscillation of the solid body the hydrodynamic forces acting upon it may be divided into inertial and damping forces. The coefficients of the inertial

LC

40793

USSR/Physics (Contd)

Aug 1946

forces may be called connected masses, this being a generalization of the existing concept of a connected mass for an infinite and incompressible liquid. The damping forces account for the continuous expenditure of energy on the formation of acoustical waves and are linearly dependent upon velocities. The same properties of symmetry hold true as well for the coefficients of damping as for the generalization of the connected masses.

IC

40793

[illegible][illegible]

51

一、

Leborch, S. V. and Maslov, V.
of finite span in a sequence, 24
and New York State, 1964
and New York State, 1964
and New York State, 1964
and New York State, 1964
and New York State, 1964
and New York State, 1964

analytical reviews, 1964, 1965

24
F. E. M.

KHASKIND, M. D.

PA 34T101

USSR/Physics

Sound - Transmission
Velocity, Ultrasonic
Velocity, Subsonic

Jun 1947

"Transmission of Sound Through a Hole in a Gas Flow-
ing at Subsonic and Ultrasonic Velocities," M. D.
Khaskind, 6 pp

"Zhur Takh Fiz" Vol XVII, No 6, 693-698

A method is set forth for the exact solution of the
problem of the transmission of horizontal sound waves
through a hole in gas which is moving at subsonic and
ultrasonic speed. The examination of this simple
problem makes it possible to start from the usual

USSR/Physics (Contd)

34T101
Jun 1947

equation of sound for a corresponding form of medium,
which equation is then modified in accordance with
the mobility of the medium.

34T101

KHASKIND, M. D.

"Methods of Hydrodynamics in the Problems of the Seagoing Ability of Ships
in Waves," Tr. TsAGI, No.603, 1947

KHASKIND, M.D.

I

"Davlenie Voln Na Zagranichenie," Izvestiya 1985, V. 4, No. 2, 1985.

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

Hashind, M. D., and Falkovich, S. I. Investigation of finite span in a supersonic flow

Ats. Journal of Aerodynamics, 1957, 21, 1, 1-10

Translated from Aerodinamika, 1957, 21, 1, 1-10

For AS, 1957, 21, 1, 1-10

at the Reviewer's

702

HASKIND, M. D.

①

Haskind, M. D. Oscillations of a floating contour on the surface of a heavy liquid. Akad. Nauk SSSR. Prikl. Mat. Meh. 17, 165-178 (1953). (Russian)

The author treats the oscillatory motion of a long cylindrical body (width $2a$ at the waterline) floating freely in an infinitely deep inviscid fluid; the problem is linearized. The mathematical problem is to find a harmonic function $\Phi(x, y, t) = \varphi(x, y)e^{-i\omega t}$ such that: (1) $\varphi_x = \varphi_y = 0$ ($x = \sigma^2, y$) for $y=0, |x| > a$; (2) $\varphi_x = v_x$ on the contour of the cylinder, and (3) φ has the asymptotic values $(A_1 + B_1)e^{-i\omega(x+a)}$ as $x \rightarrow +\infty$ and $(A_2 + B_2)e^{-i\omega(x-a)}$ as $x \rightarrow -\infty$. For a wide class of body profiles formulas for the force and moment on the body are derived. J. V. Wehausen.

Mathematical Reviews
Vol. 15 No. 4
Apr. 1954
Mechanics